

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to a person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment

PRECAUTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE, USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. NOTE TO PHYSICIAN: Emergency information: Call CHEMTREC (800) 424-9300

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water. Do not contaminate water when disposing of equipment washwater. Apply this product only as specified on the label.

GENERAL INFORMATION

FIRST CHOICE CROP OIL CONCENTRATE is a blend of a surfactant and a nonphytotoxic superior type of agricultural spray oil designed for use with a broad range of post-emergence herbicides as well as desiccants, defoliants and other pesticide uses. It is most beneficial with emulsion formulations of organic compounds, and is compatible with most types of herbicides, defoliants and desiccants. Avoid use with selective herbicides where the selectivity depends on not wetting the primary crop. The blend of nonionic surfactant and oil is formulated to give fast wetting and improved foliage penetration without causing premature desiccation. There will be no excessive foaming created in the spray mixtures and no corrosive effect in equipment.

DIRECTION FOR USE

MIXING: Fill spray tank 1/2 full, add the desired amount of herbicide and continue filling. Add FIRST CHOICE Crop Oil Concentrate last. After filling, continue agitation. Always follow pesticide label directions, acceptable practices and advice from your licenced crop consultant.

FULL VOLUME SPRAYS: Add 4 to 32 oz. per 100 gallons as required to wet the target foliage. This will vary with the type of foliage, nozzle pressure, amount of wetting agent in herbicide, etc. and must be determined for the conditions involved.

CONCENTRATE SPRAYS: Add 4 to 32 oz. per acre as required by the type of foliage, equipment, and the herbicide wetting requirements to obtain fast, smooth wetting. Fast wetting is especially critical with air blast sprayers and with airplane application.

The following table of suggested uses is intended to serve as a guide for the use of FIRST CHOICE Crop Oil Concentrate, although the use of FIRST CHOICE Crop Oil Concentrate is not limited to these herbicides, desiccants and defoliants of these crops.

STORAGE AND DISPOSAL

STORAGE: Store in original container only. Do not contaminate water, food or feed by storage and disposal. If a spill occurs, visible puddles or drops should be covered with a dry absorbent such as clay, sawdust, lime or soda ash. Sweep up and dispose of in a permitted disposal facility. Avoid cross-contamination with other pesticides and fertilizers during handling and storage. Do not store near heat or open flame. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. In case of spill or accident, call Chemtrec (800) 424-9300.

CONTAINER DISPOSAL: Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state or local ACRC recycler, visit the ACRC web page at www.acrecycle.org. Decontaminated containers may also be disposed of in a sanitary land fill.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemically resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (such as barrier laminate, nitrile rubber, neoprene rubber, or viton).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.



**CROP OIL
CONCENTRATE**

**READ ENTIRE CONTAINER LABEL
BEFORE USING PRODUCT**

PRINCIPAL FUNCTIONING AGENTS:

Paraffin-based petroleum oil, tall oil fatty acids and alkylphenol ethoxylate.....99.0%
CONSTITUENTS INEFFECTIVE AS SPRAY ADJUVANT.....1.0%
Contains Petroleum distillates.

WA Reg. No. 11656-70004

NET CONTENTS: _____ GALLONS (_____ LITERS)

Manufactured for:
WESTERN FARM SERVICE, INC.
P.O. BOX 1168
Fresno, CA 93715

CONDITIONS OF SALE AND WARRANTY

Seller warrants that the product conforms to it's chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.

BASAGRAN, POAST, and PYRAMIN are registered trademarks of BASF Ag.
PROVADO is a registered trademark of Bayer.
SINBAR is a registered trademark of DuPont.
AGRIMEK, GRAMOXONE EXTRA & MAX, DIQUAT and FUSILADE are registered trademarks of Syngenta Crop Protection.
ROUNDUP Original, Ultra, and Weathermax are registered trademarks of Monsanto.
GOAL is a registered trademark of Dow Agro Science.
GALIGAN is a registered trademark of Makhtheshim-Agan North America.
BOA is a registered trademark of Griffin LLC.

DIRECTION FOR USE

CROPS	WEEDS	HERBICIDE	DOSAGE	REMARKS
		(Per Label Recommendations)	FC Crop Oil Concentrate/spray	
APPLES, PEACHES, CITRUS, GRAPES, CHERRIES, POME FRUITS, STONE and NUT	Annual Broadleaf and Grasses	Gramoxone Extra	GROUND 1 pint-2 quarts/ 15-40 gallons	
	Annual and Perennial Grasses and Broadleaf	Terbacil	GROUND 1 quart/40 gallons	
CORN Field or Sweet	Annual Weeds and some Perennials	Simazine	GROUND 1 quart/25 gallons	
	Annual and Perennial Grasses	Dalapon	GROUND 1 quart/50 gallons	
	Annual Broadleaf and Grasses	Atrazine	GROUND 1 quart/15 gallons AIR 1 quart/5 gallons	Apply postemergence before weeds are 1-1/2' high
		Gramoxone Extra or Atrazine	GROUND 1 pint - 1 quart/ 100 gallons	Preplant treatment for non-till corn
		Linuron	GROUND 1 quart/15-25 gallons	Directed spray only
	Giant Foxtail	Atrazine	GROUND 2 quarts/15 gallons AIR 2 quarts/5 gallons	Apply postemergence before weeds are 1-1/2' high
	Annual and Perennial Grasses and Broadleaf	Atrazine and/or Dicamba	GROUND 2 quarts/20 gallons	
	Annual and Perennial Grasses	Dalapon	GROUND 1 quart/25-50 gallons	Directed spray only
	Annual Broadleaf	2,4-D Ester and Amino Formulations	GROUND 1 quart/15-50 gallons	After silking only
PASTURE RENOVATION and BRUSH CONTROL	Annual and Perennial Grasses	Dalapon	GROUND 1-2 quarts/50 gallons	Roadways, parkways, rights-of-way, fence lines, parking lots, highway margins, storage areas, around commercial buildings, power plants, and similar areas.
	Annual Broadleaf	2,4-D Ester and Amino Formulations	GROUND 1 quart/15-50 gallons	
POTATO VINE KILLING		Gramoxone Extra Diquat	GROUND 2 quarts per acre with normal amount of water	If copper fungicides are used with killing agents, check compatibility
		Ametryne	AIR 2 quarts/5 gallons	
MINT		Poast Gramoxone Extra Sintar or Fusilade	GROUND 1 quart per acre with normal amount of water	If copper fungicides are used with killing agents, check compatibility
			AIR 1-2 quarts/5 gallon	
RICE	General	Propanil	GROUND 1 pt.-1 qt./30 gals. AIR 1 pint/5 gallons	
SEED CROPS ALFALFA, CLOVER and VETCH		Gramoxone Extra Diquat	GROUND 2-4 qts./40 gals. AIR 2-4 quarts/10 gallons	Preharvest defoliation and desiccation
LEGUMES		Poast	GROUND 1 quart per acre with normal amount of water. AIR 1 quart per acre	
SORGHUM or MILO		Same as corn recommendations		Apply after sorghum is 2 inches tall and less than 6 inches tall when weeds are less than 1-1/2 inches tall
HOPS		Gramoxone Extra	GROUND 1-6 quarts per acre with normal amount of water	
SUGAR BEETS	Annual Broadleaf and Grasses	Pyramin W with Dalapon	GROUND 1 pint/25 gallons spray	Consult your best sugar company fieldman and extension agent. Do not use FC Crop Oil Concentrate with PYRAMIN PLUS
NON-CROP MISC.	Annual and Perennial Grasses and Broadleaf	Diuron or Bromacil Simazine	GROUND 2 quarts /50 gallons GROUND 2 quarts/5 gallons	Roadways, parkways, rights-of-way, fence lines, parking lots, highway margins, storage areas, around commercial buildings, power plants, and similar areas.
	Annual Broadleaf and Grasses	MSMA	GROUND 1-2 quarts/100 gallons	
	Annual and Perennial Grasses	Dalapon	GROUND 1 pint-1 quart/50 gallons GROUND 1 quart/50 gallons	
CANOLA	Grasses		GROUND 2 quarts per acre with normal amount of water. AIR .5% of spray solution	
COTTON		Use according to Herbicide/Desiccant label recommendations and accepted practices for both weed control and to aid in harvest		

ADDITIONAL PESTICIDE USES

CROPS	PESTICIDES	DOSAGE	REMARKS
PEARS	Agrimek	Apply up to 2.5% by spray volume. One gallon per acre in final finish spray.	Do not apply with sulfur sprays.
GRAPES	Gramoxone Max, Poast, Goal, Galigan, RoundUp, and BOA	Apply 2% by spray volume.	Do not apply with sulfur sprays.
HOPS	Agrimek or Provado	Apply a maximum of .37% by spray volume per acre/1.5 quarts per acre in 100 gallon spray solution.	Do not exceed 100 gallon spray volume per acre. Do not spray within 2 weeks of Omite or Sulfur sprays.