

IN CASE OF EMERGENCY
CALL CHEMTREC: (800) 424-9300



CORROSIVE LIQUID, N.O.S.
UN 1760
(CONTAINS PHOSPHORIC ACID)
DO NOT REUSE CONTAINER

CONDITIONS OF SALE AND WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk or any such use. These risks include, but are not limited to damage to plants and crops to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

NET CONTENTS: 4 X 1 GALLONS
Weight 10.73 Lbs./gallon at 68° F

Manufactured by: Western Farm Service, Inc.
P.O. Box 1168, Fresno, CA 93715

First Choice® is a Registered Trademark of Western Farm Service, Inc.
05-02(02)dhf



Foliar Nutrient 3
10-12-0 Solution

ANALYSIS AS NUTRIENT SPRAY

Nitrogen (N)	10.0%
10% Urea Nitrogen	
Available Phosphate(P ₂ O ₅)	12.0%
Manganese (Expressed as elemental).....	1.0%
Zinc (Expressed as elemental).....	2.0%
Nitrogen derived from urea. Available phosphate derived from phosphoric acid. Manganese and Zinc derived from the sulfates.	

ANALYSIS AS SPRAY ADJUVANT:

PRINCIPAL INGREDIENTS:	
Phosphoric Acid	16.17%
Manganese Sulfate	3.20%
Zinc Sulfate	5.60%
OTHER INGREDIENTS	75.03%
TOTAL	100.00%

CA Reg. No. 11656-50102-AA

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1-2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.
IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.westernfarmsservice.com/non-nutritivemetals.com>

READ ENTIRE LABEL
USE STRICTLY IN ACCORDANCE
WITH CAUTIONS AND DIRECTIONS

PRECAUTIONARY STATEMENTS

Caution. Causes moderate eye irritation. Harmful if swallowed. Causes skin irritation. Do not get in eyes, on skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. STORAGE - Store in a dry location away from children, animals, food, feeds and seeds. Do not store near any source of heat or flame. Store only in original container. DISPOSAL - Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

ACIDIFYING-BUFFER AGENT: Will lower the pH of alkaline water thus increasing the effectiveness of all phosphate, carbamate and chlorinated hydro-carbon insecticides and lengthening their residual life. COMPATIBILITY: Compatible with most insecticides and minor elements with the exception of compounds containing sabadilla used on citrus. Caution should be used with Copper fungicides as this material may increase their solubility.

SUGGESTED USES

ALMONDS APRICOTS, PLUMS, NECTARINES, PEACHES, WALNUTS: Apply in delayed dormant period continuing through spring period. Rate: 1 to 2 quarts per acre per application. Several applications per year are advised.
ALFALFA: For seed formation apply in period prior to bloom followed by 3-4 applications every 10 days. Rate: 1 to 2 quarts per acre in 10 to 30 gallons of water.
ASPARAGUS, SPINACH, CARROTS, CELERY, GARLIC, ONIONS, PEPPERS, POTATOES, ARTICHOKEs: Apply 1 to 2 quarts per acre as often as needed.
BARLEY, WHEAT, RICE, MILO, OATS: Apply in seedling stage. 1 to 2 quarts per acre per application.
BEANS, PEAS: 1 to 2 quarts per acre just prior to bloom stage.
BOYSENBERRIES, BLACKBERRIES, STRAWBERRIES: Apply 1 to 2 quarts per acre.
CITRUS: Apply 1 to 2 quarts per acre.
COLE CROPS: Broccoli, Brussel Sprouts, Cabbage, Cauliflower: Apply 1 to 2 quarts per acre. 1 to 2 applications per season.
CORN: Apply 1 to 2 quarts to 20 gallons of water per acre. Best period is when the tassels are first appearing.
COTTON: 1 to 3 applications starting at period prior to main bloom. Dosage: 1 to 3 quarts per acre.
GRAPES: 3 applications per year, starting with period of bloom. Dosage: 1 to 2 quarts per year.
LETTUCE: 1 to 4 applications at 10-day intervals starting at thinning period. Use 1 to 2 quarts per acre.
MELONS, CANTALOUPEs, CUCUMBERS, ITALIAN SQUASH: Use 1 to 2 quarts per acre with maximum concentration 1 to 2 quarts to 15 gallons of water.
ORNAMENTALS AND FLOWER CROPS: Apply 1 to 2 quarts per acre in sufficient water to cover.
SUGAR BEETS: 1 to 2 quarts per acre. Several applications at 10 to 15 day intervals may be needed.
TOMATOES: 1 to 2 quarts per acre every 10 to 15 days in the period just before and during fruiting period.