



Super K Plus 10-10-36 plus sulfur and micronutrients

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GUARANTEED ANALYSIS

Total Nitrogen (N).....	10.0%
9.0% Nitrate Nitrogen	
1.0% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)...	10.0%
Soluble Potash (K ₂ O).....	36.0%
Sulfur (S).....	1.00%
1.00% Sulfur (S) Combined	
Boron (B).....	0.50%
0.50% Soluble Boron	
Copper (Cu).....	0.25%
0.25% Soluble Copper	
Iron (Fe).....	0.10%
0.10% Chelated Iron	
Manganese (Mn).....	0.10%
0.10% Soluble Manganese	
Molybdenum (Mo).....	0.0005%
0.0005% Soluble Molybdenum	
Zinc (Zn).....	1.00%
1.00% Soluble Zinc	

Derived from Ammonium sulfate, mono potassium phosphate, potassium nitrate, urea, sodium borate, copper sulfate, iron EDTA, manganese sulfate, sodium molybdate and zinc sulfate.

CAUTION

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
AVOID BREATHING DUST
THIS PRODUCT CONTAINS BORON.
USE OF THIS PRODUCT OTHER THAN
THAT RECOMMENDED MAY RESULT IN
INJURY TO CROPS**

GENERAL DIRECTIONS

FIRST CHOICE SUPER K PLUS is generally compatible with pesticides and other fertilizers. Determine compatibility with a jar test before mixing products

APPLICATION IN IRRIGATION WATER

Solid Set, Overhead, Sprinkler or Pivot Irrigation Systems: Dissolve FIRST CHOICE SUPER K PLUS in mix tank and apply 8-10 lbs. per acre in irrigation water. Use of anti-siphon or back-check valve is required to prevent material from running back into primary water supply.

DILUTE APPLICATIONS: 3 to 5 lbs. per 100 gal. of water.

SEMI-CONCENTRATE: 5 to 20 lbs. per acre in 40 to 100 gal. of water.

AIR APPLICATION: 5 to 10 lbs. per acre in sufficient water for good coverage.

- Fill tank with water. Start agitation. Add FIRST CHOICE SUPER K PLUS slowly and continue agitating until product is completely dissolved. Use this procedure only. DO NOT pre-mix as a slurry. Add pesticide last.
- Exceeding 2 lbs. FIRST CHOICE SUPER K PLUS, or other materials per gallon of water will over-saturate spray solution and cause a precipitate.
- When temperatures are cold, allow extra time for this product to completely dissolve.
- If additional wetting is required, add a non-ionic spreader at 1 pint per 100 gal. or 1 pint per acre by air.
- Avoid foliage application when plants are suffering from abnormal growing conditions due to heat stress or drought.

•Do not use on Golden Delicious during slow drying conditions.

SMALL GRAINS: (Wheat, oats, barley, rye). Use 5 to 10 lbs. per acre in sufficient water to cover by air or ground. Apply at tillering and again at early boot stage.

CAUTION: Use only with ester forms when combining with 2,4D. Do not use with amine form of 2,4D or dicamba (Banvel).

DECIDUOUS FRUITS: (Apples, pears, cherries, plums, prunes, peaches, almonds, walnuts, filberts, pistachios and pecans). Apply in pre-bloom and post-bloom sprays at 3 to 5 lbs. per 100 gal. in dilute sprays or 10 to 15 lbs. per acre in concentrate sprays. Avoid combining with materials that may cause russetting. Post-harvest application of 10 to 40 lbs. per acre can enhance tree vigor and increase nutrient levels with in spurs the following season.

HOPS: Ground sprayers: Use 8 to 10 lbs. per acre in 100 to 200 gallons of water. Air application use 5 to 10 lbs. per acre in no less than 10 gal. of water. Apply when hops are halfway up wire and again when they reach the top of the wire.

GRAPES: Ground sprayers: use 5 to 10 lbs. per acre in 50 to 100 gal. of water. Air application: use 5 to 10 lbs. per acre in 10 gals. of water. Apply when leaves emerge, pre-bloom and post-bloom until grapes are ¼ size.

POTATOES: Use 10 lbs. per acre. Apply when plants are 4" to 6" high (Hooking stage). Subsequent applications at 2 to 3 week intervals. Later applications may prevent "early die" symptoms.

CORN: (Field and Sweet) Use 8 to 10 lbs. per acre. Apply when corn is 6" to 8" high and again prior to tasseling.

BEANS, PEAS, LENTILS, RAPESEED (Canola): Use 5 to 10 lbs. per acre. Apply at first sign of bloom. Repeat 2 weeks later.

COLE CROPS, LETTUCE, LEAFY VEGETABLES: Use 8 to 10 lbs. per acre. Apply right after thinning or transplanting. Repeat every 10 to 14 days. Additional wetting may be needed.

SUGAR BEETS: Use 8 to 10 lbs. per acre. Apply when leaves are 10" across. Repeat 3 to 4 weeks later. Additional wetting may be needed.

FIELD CROPS: (mint, alfalfa, clover) Use 8 to 10 lbs. per acre. Apply when plants have 6" to 8" new growth. Repeat to new growth after each cutting.

ONIONS, GARLIC, TOMATOES, PEPPERS: Use 8 to 10 lbs. per acre. Begin application prior to bloom. Repeat every 2 to 3 weeks thereafter.

BERRIES: (Strawberries, raspberries, blackberries, blueberries) Use 5 to 10 lbs. per acre in spring on new growth. Repeat at pre-bloom and 3 weeks later. Use sufficient water for good coverage.

CRANBERRIES: Use 8 to 10 lbs. per acre. Apply a popcorn stage. Repeat at hooking and 3 weeks later

GARDENS, SHRUBS, FLOWERS, LAWNS: Dissolve ½ lb. In 10 gal. water and apply to foliage with sprayer. For ground application, use 1 lb. per 1,000 square feet.

COMPATIBILITY: FIRST CHOICE SUPER K PLUS is compatible with most commonly used pesticides. Do not use with lead arsenate, hydrated lime, lime sulfur or other strongly alkaline materials. Do not use with amine form of 2,4D or with dicamba (e.g. Banvel). In all cases, it is recommended that a jar test be made to determine compatibility of FIRST CHOICE SUPER K with other materials before proceeding with high volume mixing.

CONDITIONS OF SALE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured For:
Western Farm Service, Inc.
Jerome, Idaho 83338

NET WEIGHT 50 POUNDS (22.68 Kg.)

First Choice® is a registered trademark of Western Farm Service, Inc.