



Bollster 10-0-9

Plus 9% Ca-1%B-0.2%Zn

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GUARANTEED ANALYSIS:

Total Nitrogen (N).....	10.00%
0.45% Ammonical Nitrogen	
9.35% Nitrate Nitrogen	
0.2% Urea Nitrogen	
Soluble Potash (K ₂ O).....	9.00%
Calcium (Ca)	9.00%
Boron (B)	1.00%
Zinc (Zn)	0.20%
0.20 Chelated Zinc	

Derived from:
Calcium Nitrate, Calcium Ammonium Nitrate, Potassium Nitrate, Urea, Sodium Borate and Zinc EDTA.

CONDITIONS OF SALE

1. Seller warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Seller neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond seller's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein, and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by seller under 1. above) resulting from handling, storage and use of the material.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Excess amounts of Boron (B) can cause plant injury.

NET CONTENTS: 2½ GALLONS / 9.463 LITERS
13.35 lbs./gallon

Manufactured for:
Western Farm Service, Inc.
P. O. Box 1168
Fresno California 93715

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.

GENERAL INFORMATION

FIRST CHOICE Bollster is designed to enhance growth and yield. FIRST CHOICE Bollster may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. FIRST CHOICE Bollster is recommended as foliar nutrition to be applied to cotton where peak nutrient demand is indicated by laboratory tissue analysis.

DIRECTIONS FOR USE

Dissolve the desired amount of FIRST CHOICE Bollster in water before adding to the spray tank. With proper agitation running, add FIRST CHOICE Bollster to the spray tank. Add FIRST CHOICE Bollster to the spray tank before adding pesticides.

Spray immediately after mixing. Do not store solution mixed with pesticide!

SUGGESTED USE RATES:

CROP	RATE/ACRE	NUMBER & TIMING OF APPLICATIONS
Cotton	3-6 pts/acre	3-4 applications per season. Time applications at pre bloom, 7-10 days later (10-12 nodes), and 10-14 days later (12-14 nodes). Time fourth application as necessary.
For aerial application use a maximum of two 2 quarts of FIRST CHOICE Bollster per 10 gallons of water		

Rates suggested are for standard spray volumes of 40-150 gal/acre. For higher volume sprays, rate should be adjusted.

Caution: Store FIRST CHOICE Bollster in temperatures between 41°F to 104°F and avoid extreme variations in temperature.

Use caution when applying to fruiting crops in combination with pesticides and/or surfactants. Use minimal effective rates of stickers during ripening. Do not use high analysis organo silicones or high analysis non-ionics during ripening. Avoid application to fruit at elevated temperatures (>95°F). Avoid applications to crops under environmental stress or pest pressure. Maximum effectiveness will be obtained when applied early in the morning or after dusk.