



# CalMax® Plus

## Low Chloride Suspension Calcium

### 10-5-3

**READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT**

**GUARANTEED ANALYSIS:**

Total Nitrogen (N).....	10.0%
0.58% Ammoniacal Nitrogen	
8.10% Nitrate Nitrogen	
1.32% Urea Nitrogen	
Available Phosphoric Acid(P <sub>2</sub> O <sub>5</sub> ).....	5.0%
Soluble Potash (K <sub>2</sub> O).....	3.0%
Calcium (Ca).....	11.0%
Boron (B).....	0.05%
Copper (Cu).....	0.05%
0.05% Chelated Copper	
Iron (Fe).....	0.10%
0.10% Chelated Iron	
Manganese (Mn).....	0.10%
0.10% Chelated Manganese	
Molybdenum (Mo).....	0.001%
Zinc (Zn).....	0.15%
0.15% Chelated Zinc	

Derived from Calcium Nitrate, Calcium Ammonium Nitrate, Urea, Monopotassium Phosphate, Boric Acid, Copper EDTA, Iron EDTA, Manganese EDTA, Sodium Molybdate, Zinc EDTA.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.  
 WARNING: Excess amount of Boron (B) can cause plant injury.

**This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.**

**GENERAL INFORMATION**

FIRST CHOICE CalMax Plus is a water soluble foliar nutritional fertilizer of macro- and micro nutrients. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended.

Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. FIRST CHOICE CalMax Plus is recommended as foliar nutrition to be applied to crops where disorders due to insufficient calcium are anticipated.

**CONDITIONS OF SALE AND WARRANTY**

1. Seller warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Seller neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond seller's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon, and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by seller under 1. above) resulting from handling, storage and use of the material.

NET CONTENTS: 2.5 GALLONS(9.46 LITERS)  
 12.52 lbs./gallon(1.50 kgs./liter)

Manufactured for:  
**Western Farm Service, Inc.**  
 P. O. Box 1168  
 Fresno California 93715

FIRST CHOICE is a registered trademark of Western Farm Service, Inc. CalMax is a registered trademark of Omex Agrifluids.

## DIRECTIONS FOR USE

Dissolve the desired amount of FIRST CHOICE CalMax Plus in water before adding to the spray tank. With proper agitation, FIRST CHOICE CalMax Plus can be added into the spray tank with agitation running. Do not store solution mixed with pesticide! Spray immediately after mixing. Add FIRST CHOICE CalMax Plus to the spray tank before adding pesticides.

### RECOMMENDATIONS FOR USE:

CROP	PURPOSE	RATE/ACRE	RATE/HA	NUMBER & TIMING OF APPLICATIONS
Apples	Bitter pit reduction	3-4 pints	3.5-5.0 liters	5 applications starting at petal fall. Combine with cover sprays.
Avocados	Pulp spot reduction	3.2-6.8 pints	4.0-8.5 liters	Multiple applications
Cabbage, Cauliflower Lettuce, Endive	Tip burn reduction	3-4 pints	3.5-5.0 liters	4-6 applications starting shortly before head formation.
Chinese Cabbage	Tip burn reduction	2-4 pints	2.5-5.0 liters	2-4 applications per season
Broccoli	Brown bead reduction	3-4 pints	3.5-5.0 liters	4-6 applications starting shortly before bead formation.
Brussels Sprouts	Internal browning reduction	3.2-6.8 pints	4.0-8.5 liters	Multiple applications
Celery, Chicory	Black heart reduction	3-4 pints	3.5-5.0 liters	Weekly applications starting before black heart symptoms arise.
Cherries, Plums	Cracking reduction	3-5 pints	3.5-6.0 liters	3-4 applications starting 6-8 weeks before harvest.
Cotton	Square shedding reduction	5 pints	6.0 liters	3 applications between 5-7 leaf stage and flowering.
Cucumbers, Melons, Peppers, Tomatoes	Blossom end rot reduction	1-3 pints	1.5-3.5 liters	6-12 applications during fruit development.
Potatoes	Internal brown spot reduction	2-4 pints	2.5-5.0 liters	Multiple applications during periods of heat stress.
Strawberries and other berries	Increased fruit firmness	5 pints	6.0 liters	3 applications in conjunction with last pre-harvest pesticide sprays.
Grapes	Reduction of stem dieback and shot berry	4-8 pints	5.0-10.0 liters	3-4 applications from beginning of berry softening to maturity
Table Grapes	Improved fruit finish and storability	4-5 pints	5.0-6.0 liters	
Kiwi	Blossom end rot reduction	3.2-6.8 pints	4.0-8.5 liters	Multiple applications
Pears	Superficial scald reduction	3.2-6.8 pints	4.0-8.5 liters	Multiple applications
Peaches Nectarines	Improved fruit firmness	3-4 pints	3.5-5.0 liters	4-5 treatments beginning from fruit-set
Ornamentals: Poinsettia	Reduction of marginal bract necrosis	Concentration 0.4-0.6%		2-4 Applications weekly, stopping at the stage of bract colorations
For aerial application use a maximum of two (2) quarts of FIRST CHOICE CalMax Plus per 10 gallons of water.				

Rates recommended are for standard spray volumes of 50-150 gal/acre (500-1,500 liters/ha.). For higher volume sprays, rate should be adjusted. Caution: Store FIRST CHOICE CalMax Plus in temperatures between 41 F to 104 F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after long storage, this however does not affect quality or effectiveness of the product.