



Foliar Advance

FOLIAR NUTRITION FOR VITAL PLANT GROWTH STAGES

2-10-10

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GUARANTEED ANALYSIS 2-10-10

Nitrogen (N).....	2.00%
1.0% Ammonical Nitrogen	
1.0% Urea Nitrogen	
Available Phosphate (P ₂ O ₅).....	10.00%
Soluble Potash (K ₂ O).....	10.00%
Boron (B)	0.02%
Molybdenum (Mo)	0.001%

Derived from urea, dipotassium phosphate, boric acid and sodium molybdate.

ALSO CONTAINS NON PLANT FOOD INGREDIENTS

0.015% Humic Acid derived from leonardite.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes since it causes irritation and possible damage. Thoroughly flush any exposed body part with liberal quantities of water. Get medical attention for eyes.

DIRECTIONS FOR USE SHAKE WELL BEFORE USING

DO NOT EXCEED 3 gallons per 10 gallons of finished spray.

APPLICATION – Thorough coverage and wetting is desirable for optimum foliar spray results.

CAUTION – DO NOT use with oil, lime sulfur or highly alkaline materials. Conduct a jar test before tank mixing new or untested mixture to determine compatibility.

APPLICATION – AIR – Apply in 10 gallons of water per acre – **GROUND** – apply in 20 to 200 gallons of water per acre.

FILBERT OR HAZEL NUTS

NUT BEARING ORCHARDS – Apply 1.5 to 3 gallons per acre. Use higher rates where the tree canopy is closed for best results. Apply one

application after nut harvest prior to November 15. **DO NOT** apply if air temperatures are expected to drop below 20 degrees Fahrenheit within 14 days following application

NON-BEARING AND YOUNG ORCHARDS-

(less than 5 years old) – Apply 0.75 to 1.5 gallons to the trees in the spring after two to three leaves have fully emerged and make a second application of 0.75 to 1.5 gallons per acre in the fall prior to November 15.

NURSERIES

NON-BEARING FRUIT TREES – Apply 6 gallons per acre or two 3 gallons per acre applications as directed spray to the trees when the trees have reached the stage or height where increase lateral branching is desired.

BEARING FRUIT TREES

SWEET CHERRIES – Apply 2 applications of 1.5 gallons per acre. The first application in the fall (September thru October) and a second application of 1.5 gallons per acre in the spring (at blossom bud swell or popcorn stage).

CONIFER SEEDLINGS AND SEEDLING

CHRISTMAS TREES – Apply 2 applications per season. Apply the first application of 1.5 gallons per acre to the primary needles or newly emerged seedlings or just prior to first bud swell in the spring of older seedlings. For best results, a second application of 1.5 gallons per acre should be applied just prior to bud break in conifers when a second flush occurs.

FIELD AND VEGETABLE CROPS

ALFALFA HAY – Apply 1.5 gallons per acre within 1-7 days following swathing of each cutting.

CORN (SWEET AND FIELD) – Apply 1-2 applications of 1.5 gallons per acre, beginning at the 3-6 leaf stage. The second application of 1.5 gallons should be made 14-21 days later. **DO NOT ADD SURFACTANT** when applying to corn. **GARLIC** – Apply 1-2 applications of 1.5 gallons per acre. Apply the first application 2 weeks after emergence and the second application 2-3 weeks later.

ONIONS – Apply 1-2 applications 1.5 gallons per acre. Apply the first application 2-3 leaves and a

second application when the onions have 4-5 leaves.

POTATOES – Apply a MINIMUM OF TWO APPLICATIONS at the rate of 1.5 gallons per acre during the season for best results. Apply the first application when the potatoes begin to hook (form tubers) and the second application 7-21 days later.

RED CLOVER FOR SEED – Apply 1.5 gallons per acre within 1-5 days following green chopping in the spring.

BERRIES

RASPBERRIES AND BLACKBERRIES – Apply 2 applications of 1.5 gallons per acre beginning in the later summer (September thru October) directing the spray solution to the next seasons fruiting canes. Apply a second application of 1.5 gallons per acre in the late winter-early spring (mid February thru March) but just prior to bud break.

BLUEBERRIES – Apply 1-2 applications of 1.5 gallons per acre. The first application in the fall (September thru October) and a second application of 1.5 gallons per acre in the early spring (March-April) early blossom bud swelling to pollination.

CONDITIONS OF SALE

1. Seller warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Seller neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond seller's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein, and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by seller under 1. above) resulting from handling, storage and use of the material.

NET CONTENTS:
30 Gallons
10 lbs. / gallon

Manufactured by:
Western Farm Service, Inc.
P. O. Box 1168
Fresno California 93715

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.

06-02(02) dhf

