



Foliar Advantage K

Low Chloride Foliar Suspension

2-0-27

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GUARANTEED ANALYSIS:

Total Nitrogen (N).....2.00%
2.00% Urea Nitrogen
Soluble Potash (K₂O)27.00%
Magnesium (Mg)1.00%
Sulfur (S)10.70%

Derived from: Potassium Sulfate,
Magnesium Sulfate and Urea.

KEEP OUT OF REACH OF CHILDREN
CAUTION

GENERAL INFORMATION

FIRST CHOICE Foliar Advantage K is a water-soluble foliar nutritional fertilizer of macro and micronutrients. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. FIRST CHOICE Foliar Advantage K is recommended as foliar nutrition to be applied to crops where insufficient nutrient levels are indicated by laboratory tissue analysis.

CONDITIONS OF SALE

1. Seller warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Seller neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond seller's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein, and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by seller under 1. above) resulting from handling, storage and use of the material.

NET CONTENTS:

2½ GALLONS / 9.463 LITERS
12.10 lbs./gallon

Manufactured for:
Western Farm Service, Inc.
P. O. Box 1168
Fresno California 93715

DIRECTIONS FOR USE

Dissolve the desired amount of FIRST CHOICE Foliar Advantage K in water before adding to the spray tank. With proper agitation running, add FIRST CHOICE Foliar Advantage K to the spray tank. Add FIRST CHOICE Foliar Advantage K to the spray tank before adding pesticides. Spray immediately after mixing. Do not store solution mixed with pesticide!

SUGGESTED USE RATES:

CROP	RATE/ACRE	NUMBER & TIMING OF APPLICATIONS
Alfalfa	3-6 pts/acre	Apply after second and fifth cuttings at 4"-6" of re-growth.
Almonds, Pistachios	3-6 pts/acre	2-3 applications as required from jacket split to hull split as required.
Apples	3-6 pts/acre	2-3 applications starting at post thinning through early sizing, including 14-21 days post harvest.
Apricots, Nectarines, Peaches	3-4 pts/acre	2-3 applications from shuck split to full color, including 14-21 days post harvest.
Avocados	3-6 pts/acre	Multiple applications as required. Do not apply within 21 days of harvest.
Caneberries	3-4 pts/acre	2 applications, one each immediately post bloom and 21-30 days later.
Celery, Chicory	2-4 pts/acre	2-3 applications during the season as required.
Cherries, Plums, Prunes	3-6 pts/acre	2-3 applications starting at jacket split to full color, including post harvest.
Citrus	3-6 pts/acre	Multiple applications as required.
Cotton	3-5 pts/acre	2-3 applications from squaring to full boll development.
Cucumbers, Melons	3-4 pts/acre	2-3 applications from full bloom through fruit sizing.
Dry Beans such as Kidneys, Pintos, Limas	3-4 pts/acre	2 applications, one each immediately post bloom and again 30-45 days later.
Grapes	2-4 pts/acre	2-3 applications from post shatter through early ripening.
Kiwi	3-4 pts/acre	2-3 applications from early fruit formation to ripening.
Ornamentals: Woody Plants	2-4 pts/acre	Multiple applications as required during growing season with one application after September 1 st for winter hardiness.
Pears	3-6 pts/acre	3 applications, one each at early fruit formation, 30 days pre-harvest, and 14-21 days post harvest.
Peppers, Tomatoes	3-5 pts/acre	2-3 applications from fruit sizing to first color.
Potatoes	3-4 pts/acre	2-3 applications from flowering through tuberization.
Strawberries	2-4 pts/acre	2 applications, one each immediately post bloom and 21-30 days later.
Sugar Beets	3-4 pts/acre	2-3 applications from early beet development at 21-30 day intervals.
Sweet Potatoes	3-4 pts/acre	2-3 applications from flowering through tuberization.
Table Grapes	3-4 pts/acre	2-3 applications as required from early fruit formation to ripening.
Walnuts	3-6 pts/acre	2-3 applications from sizing through packing tissue browning.
For aerial application use a maximum of 3 pints of FIRST CHOICE Foliar Advantage K per 10 gallons of water.		

Rates suggested are for standard spray volumes of 40-150 gal/acre. For higher volume sprays, rate should be adjusted. Caution: Store FIRST CHOICE Foliar Advantage K in temperatures between 41°F to 104°F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after long storage, this however does not affect quality or effectiveness of the product.

Use caution when applying to fruiting crops in combination with pesticides and/or surfactants. Use minimal effective rates of stickers during ripening. Do not use high analysis organo silicones or high analysis non-ionics during ripening. Avoid application to fruit at elevated temperatures(>95°F). Avoid applications to crops under environmental stress or pest pressure. Maximum effectiveness will be obtained when applied early in the morning or after dusk.