



# FOLIAR EXTRA

## 6-12-7

### GUARANTEED ANALYSIS

Total Nitrogen (N) .....	6%
1.5% Ammoniacal Nitrogen	
4.5% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	12%
Soluble Potash (K <sub>2</sub> O) .....	7%
Weight 10.38 lbs. per gallon	

Derived from urea and ammonium phosphate, phosphoric acid and potassium hydroxide.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS

0.20% humic acid derived from leonardite.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

**WARNING:** This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

### CONDITIONS OF SALE AND WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, damage caused by drift to other plants or crops, and personal injury.

**READ ENTIRE LABEL  
USE STRICTLY IN ACCORDANCE  
WITH CAUTIONS AND DIRECTIONS**

### SHAKE WELL BEFORE USING DIRECTIONS FOR USE

#### ROW CROPS

Application - Air - Apply in 5 to 10 gallons of water per acre.

Ground - Apply in 20 to 40 gallons of water per acre.

#### APPLY A MINIMUM OF TWO APPLICATIONS

BEANS - Apply 1 gallon per acre at third trifoliate leaf, make a second application of 1 gallon per acre 10-14 days later. A third application of 1 gallon per acre may be made 10-14 days after the second application.

TOMATOES - Apply 1 gallon per acre at early bloom, make a second 1 gallon application 7-10 days later. A third application of 1 gallon per acre may be made 7-10 days following the second application.

TOMATOES (Transplants) - Apply 1 gallon per acre at early bloom after transplants have begun vigorous growth, make a second application of 1 gallon per acre 7-10 days following the first. A third application of 1 gallon per acre may be made 7-10 days following the second application.

COTTON - Apply 1 gallon per acre at early bloom (5-6 blooms per 25 foot of row), make a second application of 1 gallon per acre 7-10 days following the first. A third application of 1 gallon per acre may be made 7-10 days following the second application.

STRAWBERRIES - Apply 1/2 gallon to 2 gallons per acre when first fruit buds show in the crown. Repeat at 1 to 2 week intervals for two additional applications.

DECIDUOUS FRUITS - Application-Air-Apply in 10-20 gallons of water per acre.

Ground-Apply in 100-200 gallons of water per acre. Apply 1 1/2 - 2 gallons per acre at "bud swell", make a second application of 1 1/2 - 2 gallons per acre 7-10 days following the first (at full bloom).

MELONS - Apply 1 gallon per acre at early bloom, make a second 1 gallon per acre application 7-10 days following the first. A third application of 1 gallon per acre may be made 7-10 days following the second application.

POTATOES - Apply 1 to 2 gallons per acre on the seed pieces at planting. Apply 1 gallon per acre foliar at the "hook stage", make a second foliar application of 1 gallon per acre 7-10 days following the first foliar application.

ALMONDS - Application-Air-Apply in 10-20 gallons of water per acre.

Ground - Apply in 100-200 gallons of water per acre.

#### APPLY A MINIMUM OF TWO APPLICATIONS

Apply 1.5-2 gallons per acre at "bud swell", make a second application of 1.5-2 gallons per acre 7-10 days following the first (at first bloom).

GRAPES - Application - Ground - Apply in 50 gallons of water per acre.

#### APPLY A MINIMUM OF THREE APPLICATIONS

Apply 1 gallon per acre at "bud swell", make a second application of 1 gallon per acre 7-10 days following the first, make a third application of 1 gallon per acre 7-10 days following the second application. Do not apply within 60 days of harvest or after early fruit set (pea-berry size).

COMPATIBILITY - Compatible with most pesticides and fertilizers. A compatibility test is recommended if the desired combination has not been previously used. Add pesticide after all of the water has been added.

DO NOT EXCEED - One quart per gallon of finished spray.

APPLICATION - Thorough coverage and wetting is desirable for optimum foliar spray results. Do not over spray causing runoff.

CAUTION - DO NOT use with copper sprays, oil, lime sulfur, spray lime or highly alkaline materials.

Manufactured By:

Western Farm Service, Inc.,  
P.O. Box 1168, Fresno, CA 93715

NET CONTENTS: \_\_\_\_\_ GALLONS (\_\_\_\_\_ liters)

Weight per gallon 10.8 lbs. (1.296 kgs/liter)