

FOR USE IN ID & WA ONLY



FOLIAR NUTRIENT

10-12-0

Solution

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GUARANTEED ANALYSIS:

Total Nitrogen (N)10.0%
Available Phosphoric Acid (P₂O₅)12.0%
Manganese (Mn)1.0%
Zinc (Zn)2.0%
Derived from urea, phosphoric acid, manganese sulfate and zinc sulfate.

NET WEIGHT: 10.54 lbs./gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION
HARMFUL IF SWALLOWED
AVOID EXCESSIVE SKIN CONTACT

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with a gentle stream of water for 30 minutes. Seek immediate medical attention.
IF ON SKIN: Flood contaminated area with water. Remove contaminated clothing. Seek medical attention if irritation persists.
IF SWALLOWED: Give alert patient milk or water freely. Never give anything by mouth to an unconscious person. If any unusual symptoms develop, seek immediate medical attention.
IF INHALED: Remove victim to fresh air and get medical attention if symptoms persist.

Information regarding the contents and levels of metals in this product is available on the internet at <http://agr.wa.gov/>

DIRECTIONS FOR USE

ALMONDS, APPLES, APRICOTS, PLUMS, NECTARINES, PEACHES AND WALNUTS: Apply in delayed dormant continuing through spring period. Dosage: 1 to 2 quarts per acre per application. Several applications per year advised.
ALFALFA: For seed formation apply in period prior to bloom followed by 3-4 applications every 10 days. Dosage: 1 to 2 quarts per acre in 10 to 30 gallons of water.
ASPARAGUS, SPINACH, CARROTS, CELERY, GARLIC, ONIONS, PEPPERS, POTATOES AND ARTICHOKEs: Apply 1 to 2 quarts per acre as often as needed.
BARLEY, WHEAT, RICE, MILO AND OATS: Apply in seedling stage. 1 to 2 quarts per acre per application.
BEANS AND PEAS: Apply 1 to 2 quarts per acre just prior to bloom stage.
BOYSENBERRIES, BLACKBERRIES AND STRAWBERRIES: Dosage: 1 to 2 quarts per acre.

CITRUS: 1 to 2 quarts per acre.
COLE CROPS: Broccoli, Brussel Sprouts, Cabbage and Cauliflower: 1 to 2 quarts to the acre for 1 to 2 applications.
CORN: 1 to 2 quarts to 20 gallons of water per acre. Best period is when the tassels are first appearing.
COTTON: 1 to 3 applications starting at period prior to main bloom. Dosage: 1 to 3 quarts per acre.
GRAPES: 3 applications per year, starting with period of bloom. Dosage: 1 to 2 quarts per acre.
LETTUCE: 1 to 4 applications at 10-day intervals starting at thinning period. Use 1 to 2 quarts per acre.
MELONS, CANTALOUPEs, CUCUMBERS AND ITALIAN SQUASH: Use 1 to 2 quarts per acre with maximum concentration 1 to 2 quarts to 15 gallons of water.
ORNAMENTALS AND FLOWER CROPS: Apply 1 to 2 quarts per acre in sufficient water to cover.
SUGAR BEETS: Apply 1 to 2 quarts per acre. Several applications at 10 to 15 day intervals may be justified.
TOMATOES: 1 to 2 quarts per acre every 10 to 15 days in the period just before and during fruiting period.

DO NOT REUSE CONTAINER

SHIPPING DESCRIPTION: RQ Corrosive Liquid, N.O.S. contains phosphoric acid, 8, UN1760, PG III, ERG154, Placard Corrosive-1760, Liquid Fertilizer

CONDITIONS OF SALE AND WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, damage caused by drift to other plants or crops, and personal injury.

NET CONTENTS: _____ GALLONS

Manufactured by:
WESTERN FARM SERVICE, INC.
P. O. Box 1168
Fresno CA 93715