

# FIRST CHOICE<sup>®</sup>

# GUANO PLUS<sup>™</sup>

## AGRICULTURAL MINERAL

**READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT**

**GUARANTEED ANALYSIS:**  
 Total Nitrogen (N) ..... 0.05%  
     0.015% Ammoniacal Nitrogen  
     0.035% Urea Nitrogen  
 Soluble Potash (K<sub>2</sub>O)..... 0.70%

Derived from beef and chicken compost.

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS  
 1.00% humic acid derived from leonardite.

NET WEIGHT: 8.46 lbs./gallon or 1.014 Kgs./Liter

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**This product is intended for  
 AGRICULTURAL USE ONLY**

For handling and first aid precautions please see Material  
 Safety Data Sheet (MSDS).

Information regarding the contents and levels of metals in  
 this product is available on the internet at  
<http://www.aapfco.org/metals.htm>

### DIRECTIONS FOR USE

Apply as a broadcast, banded, shanked or injected application (see directions for application). As the use of GUANO PLUS may affect crop nutrient response, timing and frequency of fertilizer applications should be based upon crop nutrition monitoring (tissue testing, post-harvest soil testing). Where warranted, adjust fertilizer rates and replace with a minimum of a 1:1 ratio of up to 15 gallons of GUANO PLUS per/acre/application. Use **20-40 gallons** per acre in a minimum of two applications during the growing season, or as needed. Split applications should apply the majority of the material in the first application (2/3 of seasonal use), and the balance during the growing season (1/3). Example: Apply 30 gallons per acre per season in a minimum of two applications. The first application (primary inoculant), of 20 gallons should be applied prior to, or at planting. Subsequent (maintenance) applications of 5-10 gallons can be made through the most convenient application means (water run, or side dress).

### DIRECTIONS FOR APPLICATION

For a seasonal total of 20-40 gpa use a combination of the following methods:

	<b>Field/row crops Gallons/Acre</b>	<b>Tree Vine Crops Gallons/Acre</b>
Broadcast	20	20
Under seed	5	--
Side dress	5-8	10
Water run/Sprinkler*	5-15	15
Micro-irrigation	10-15	5-10

\* Continue irrigation 1 hour after application to reduce risk of phytotoxicity and to facilitate product movement.

### INCOMPATIBILITIES

Do not exceed a 50 percent mix with N-P-K fertilizers or CAL PLUS unless diluted with sufficient amounts of water to ensure ease of mixing and application. Always jar test newly attempted mixes. No other known incompatibilities. Avoid strongly acidic conditions. Do not mix with Aqua Ammonia, Sulfuric Acid, Urea Sulfuric Acid or soil fumigants.

### STORAGE

Storage in a cone-bottomed tank to facilitate agitation is recommended. When stored in cone-bottomed tank, agitate 10 minutes before shipment. When stored in a flat-bottom tank, agitate 1 hour before shipment. Avoid air blast agitation, as foaming will result. If foaming does occur through normal agitation, use a quality defoamer. Agitation is recommended once every month. Agitation prior to product transfer and shipment is required. Storage for longer than 12 months is not recommended.

### HANDLING

Follow appropriate safety procedures. Skin contact may cause temporary staining. In case of accidental exposure, flush with plenty of water. Product is non-hazardous.

### SHAKE/AGITATE WELL BEFORE USING

### CONDITIONS OF SALE AND WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

NET CONTENTS: \_\_\_\_\_ GALLONS \_\_\_\_\_ LITERS

Manufactured For:  
 Western Farm Service, Inc.  
 P.O. Box 1168  
 Fresno, CA 93715

GUANO PLUS is a registered trademark of BioScientific, Inc.

06/04(02) CA-dhf