



# Hop-Phite

## 1-26-20

### Nutrients For Soil and Foliar Application

**READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT**

**GUARANTEED ANALYSIS: 1-26-20**

- Total Nitrogen (N).....1.0%
- 1.0% Ammoniacal Nitrogen
- Available Phosphoric Acid(P<sub>2</sub>O<sub>5</sub>).....26.0%
- Soluble Potash (K<sub>2</sub>O).....20.0%

Derived from Potassium polyphosphate and Ammonium polyphosphate.

**CAUTION:** Do not apply together with copper containing products.

**KEEP OUT OF REACH OF CHILDREN**

### PREVENT AND CORRECT NUTRIENT DEFICIENCIES

**GENERAL INFORMATION**

FIRST CHOICE Hop-Phite 1-26-20 is a nutritional supplement for a variety of crops. When applied properly it may be beneficial to plant growth, crop yield and quality. FIRST CHOICE Hop-Phite 1-26-20 can be used for soil and/or foliar applications. It can also be applied through irrigation water.

**DIRECTIONS FOR USE**

**SOIL APPLICATION:** FIRST CHOICE Hop-Phite 1-26-20 can be applied to soil through various types of irrigation systems. The application rate is 64 to 80 oz. per acre (5-6 liters/hectare).

**FOILAR APPLICATION:** FIRST CHOICE Hop-Phite 1-26-20 may be applied by air and concentrate or dilute ground sprayers.

ALMONDS, WALNUTS, PISTACHIOS and other Nut Crops: Apply 32-64 oz. per acre (3-5 liters/hectare) pre-bloom with subsequent applications at 30 to 90 day intervals or as needed.

APPLES, PEARS and other POME FRUITS: Apply 32-64 oz. per acre (3-5 liters/hectare) pre-bloom with subsequent applications at 30 to 90 day intervals or as needed.

APRICOTS, PEACHES and other STONE FRUITS: Apply 32-64 oz. per acre (3-5 liters/hectare) pre-bloom with subsequent applications at 30 to 90 day intervals or as needed.

AVOCADO: Apply 32-64 oz. per acre (3-5 liters/hectare) pre-bloom with subsequent applications at 30 to 90 day intervals or as needed.

BEANS and PEAS: Apply 32-64 oz. per acre (3-5 liters/hectare) after transplanting or thinning. Repeat the applications 1 to 2 times with an interval of 10 to 15 days or as needed.

BROCCOLI, CABBAGE, CAULIFLOWER, LETTUCE and SPINACH: Apply 32-64 oz. per acre (3-5 liters/hectare) after transplanting or thinning. Repeat application 1 to 2 times with an interval of 10 to 15 days or as needed.

CARROTS, BEETS, ONIONS and GARLIC: Apply 32-64 oz. per acre (3-5 liters/hectare) after transplanting or thinning. Repeat application 1 to 2 times with an interval of 10 to 15 days or as needed.

CITRUS, GRAPES, HOPS, KIWI, OLIVES: Apply 32-64 oz. per acre (3-5 liters/hectare) pre-bloom with subsequent applications at 30 to 90 day intervals or as needed.

MELONS and CUCUMBERS: Apply 32-64 oz. per acre (3-5 liters/hectare) after transplanting or thinning. Repeat application 1 to 2 times with an interval of 10 to 15 days or as needed.

POTATO and SWEET POTATO: Apply 32-64 oz. per acre (3-5 liters/hectare) when plants are 4-6 inches in height. Repeat application 1-3 times with an interval of 10 to 15 days or as needed.

RASPBERRIES and BLACKBERRIES: Apply 32-64 oz. per acre (3-5 liters/hectare) pre-bloom with subsequent applications at 30 to 90 day intervals or as needed.

STRAWBERRIES: Apply 32-64 oz. per acre (3-5 liters/hectare) after transplanting or thinning. Repeat application 1 to 2 times with an interval of the 10 to 15 days or as needed.

TOMATOES and PEPPERS: Apply 32-64 oz. per acre (3-5 liters/hectare) after transplanting or thinning. Repeat application 1 to 2 times with an interval of 10 to 15 days or as needed.

COMMERCIAL NURSERY and ORNAMENTAL CROPS: Apply 24-48 oz. per acre (2 to 4 liters/hectare) after potting or transplanting. Repeat application every 30-40 days or as needed.

*Avoid the application of copper containing products 20 days prior to the application of FIRST CHOICE*

*Hop-Phite and for 10 days after the application of  
FIRST CHOICE Hop-Phite.*

**CONDITIONS OF SALE**

1. Seller warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Seller neither makes nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond seller's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein, and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by seller under 1. above) resulting from handling, storage and use of the material.

**NET CONTENTS:**

- 1 gallon (3.78 liters)
- 2 ½ gallons (9.45 liters)
- 2 x 2 ½ gallons (18.93 liters)
- 5 gallons (18.93 liters)
- 55 gallon (207.9 liters)

**Manufactured for:  
Western Farm Service, Inc.  
P. O. Box 1168  
Fresno California 93715**

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.

02-24-00(01)-dhf