



CALMAX PK

9-9-9

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GUARANTEED ANALYSIS:

Total Nitrogen (N)	9.0%
0.45% Ammoniacal Nitrogen	
7.75% Nitrate Nitrogen	
0.80% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅).....	9.0%
Soluble Potash (K ₂ O).....	9.0%
Calcium (Ca).....	9.0%

Derived from Calcium nitrate, calcium ammonium nitrate, potassium nitrate, urea, and monopotassium phosphate.

CAUTION
Keep Out of Reach of Children

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

Information regarding the contents and levels of metals in this product is available on the internet at <http://agr.wa.gov/>

GENERAL INFORMATION

FIRST CHOICE CALMAX PK is a unique water-soluble foliar nutritional fertilizer of macronutrients and calcium. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. FIRST CHOICE CALMAX PK is recommended as a foliar nutrient to be applied to crops where insufficient nutrient levels are indicated by laboratory tissue analysis.

Use caution when applying to fruiting crops in combination with pesticides and/or surfactants. Use minimal effective rates of stickers during ripening. Do not use high analysis organo silicone or high analysis non-ionic surfactants. Avoid application to fruit at elevated temperatures (>95°F). Avoid applications to crops under environmental stress or pest pressure. Maximum effectiveness will be obtained when applied early in the morning or after dusk.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, damage caused by drift to other plants or crops, and personal injury.

NET CONTENTS: 2.5 GALLONS (9.46LITERS)
13.0 lbs./gallon(1.56 kgs./liter)

Manufactured For:
Western Farm Service, Inc.
P. O. Box 1168
Fresno California 93715

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.
CalMax® is a registered trademark of Interfarm Inc.

DIRECTIONS FOR USE

Dissolve the desired amount of FIRST CHOICE CALMAX PK in water before adding to the spray tank. With proper agitation, FIRST CHOICE CALMAX PK can be added into the spray tank with agitation running. Do not store solution mixed with pesticide! Spray immediately after mixing. Add FIRST CHOICE CALMAX PK to the spray tank before adding pesticides.

RECOMMENDATIONS FOR USE:

CROP	RATE/ACRE	NUMBER & TIMING OF APPLICATIONS
Almonds, Pistachios and Hazelnuts	4-8 pts/acre	3-4 applications starting at bud swell and repeating at 3 to 4 week intervals.
Apples	4-8 pts/acre	3-4 applications. 1 st application at bud swell, 2 nd application 1-2 weeks after petal fall, and 3-4 applications starting 6-8 weeks before harvest.
Apricots, Peaches Nectarines	4-8 pts/acre	3 applications-1 st applications at bud swell, 2 nd application at petal fall, 3 rd application at first color.
Blackberries, Blueberries Raspberries & other caneberries.	4-8 pts/acre	2 applications, one each immediately post bloom and 21-30 days later.
Brussels Sprouts	4-8 pts/acre	2-4 applications starting at thinning or transplanting & as needed to pre harvest
Celery, Chicory	4-8 pts/acre	Bi-weekly applications starting before black heart symptoms arise.
Cherries, Plums	4-8 pts/acre	1 st applications at bud swell, 2 nd application at petal fall, 3 rd application at first color and 4 th application 30-45 days pre harvest
Chinese Cabbage, cabbage, broccoli, cauliflower	4-8 pts/acre	2-4 applications per season starting at thinning or transplanting & as needed to pre harvest
Citrus & Avocados	4-8 pts/acre	Apply 1 st application prebloom. Apply subsequent applications at mid crop & 3-4 weeks prior to harvest
Cotton	4-8 pts/acre	3 applications between 5-7 leaf stage and flowering.
Cucumbers, Melons, Peppers, Tomatoes	4-8 pts/acre	2-3 applications from bloom through fruit sizing.
Head Lettuce Leaf Lettuce	4-8 pts/acre	1 st application after thinning, continue biweekly through head formation
Kiwi	4-8 pts/acre	2-3 applications starting at bud swell through early fruit sizing.
Ornamentals, conifers & Christmas trees	4-8 pints per 100 gallons	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.
Pears	4-8 pts/acre	3-4 applications. 1 st application at bud swell, 2 nd application 1-2 weeks after petal fall, and 3-4 applications starting 6-8 weeks before harvest.
Potatoes, onions & other tuber crops	4-8 pts/acre	2-3 applications starting at tuber initiation or earlier.
Strawberries	4-8 pts/acre	Apply 1 st application soon after transplanting, apply at 30 day intervals or as needed to supplement nutritional needs
Table Grapes	4-8 pts/acre	2-3 applications as required from bloom to ripening
Turf	6-10 pts/acre or 2.2-3.8 ounces per sq. ft.	Apply monthly during growing season.
Walnuts	4-8 pts/acre	2-3 applications starting a bud swell 2 nd application at catkin elongation 3 rd application at first set
Wine Grapes	4-8 pts/acre	2-3 applications starting post shatter and again at veraison

Rates recommended are for standard spray volumes of 50-150 gal/acre (500-1,500 liters/ha.). For higher volume sprays, rate should be adjusted. Caution: Store FIRST CHOICE CALMAX PK in temperatures between 41°F to 104°F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after long storage, this however does not affect quality or effectiveness of the product.