



FOLIAR GOLD

56% Sulfur Suspension

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GURANTEED ANALYSIS:

Sulfur (S)56%

Derived from: Elemental Sulfur

Contains 6.5 pounds of Sulfur (S) per gallon.

KEEP OUT REACH OF CHILDREN
Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Do not take internally. Wash thoroughly after handling.

GENERAL INFORMATION

FIRST CHOICE FOLIAR GOLD is a foliar nutritional fertilizer of macronutrients. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. FIRST CHOICE FOLIAR GOLD is recommended as foliar nutrition to be applied to crops where insufficient nutrient levels are indicated by laboratory tissue analysis.

MIXING DIRECTIONS

Add approximately 1/2 water to tank before adding FIRST CHOICE FOLIAR GOLD. Agitate thoroughly while adding remaining water. Do not allow mixture to stand. Failure to maintain agitation will cause FIRST CHOICE FOLIAR GOLD to settle and may necessitate manual stirring to redisperse. If used in combination with pesticides, add pesticides last.

COMPATABILITY

FIRST CHOICE FOLIAR GOLD is compatible with most pesticides and can be applied in existing spray programs. When using a chemical mixture that has not been used before, always try a small sample rate before application, or check compatibility by doing a jar test. Due to its acidic nature, adding FIRST CHOICE FOLIAR GOLD to the spray solution can lower the pH of the finished spray solution. Adhere to pesticide manufacturer's recommendations regarding appropriate pH range.

DIRECTIONS FOR USE

FIRST CHOICE FOLIAR GOLD can be applied to the foliage or to the soil. As a foliar application it can be applied by ground or air.

- Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality. During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures.
- Do not use FIRST CHOICE FOLIAR GOLD with oil or within 4 weeks of an oil application.
- Do not include FIRST CHOICE FOLIAR GOLD with copper sprays.
- Do not use additional surfactants without first checking with your agricultural consultant.
- Do not use when plants are under severe stress conditions such as heat or water stress or pest pressure. Avoid applications when temperatures exceed 90 degrees F.
- Use caution when applying FIRST CHOICE FOLIAR GOLD in combinations with pesticides and/or adjuvants during the fruiting stage.
- Organosilicone or high analysis non-ionic surfactants are not recommended during the ripening period.
- For maximum product effectiveness, apply early morning or after dusk.

CONDITIONS OF SALE AND WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, failure to control pests, damage caused by drift to other plants or crops, and personal injury.

NET CONTENTS: 2 1/2 GALLONS
NET WEIGHT: 11.60 lbs./gallon

Manufactured For:
WESTERN FARM SERVICE, INC.
P.O. BOX 1168
FRESNO, CA 93715

FIRST CHOICE is a registered trademark of Western Farm Service, Inc.

SUGGESTED USE RATES (foliar application):

CROP	RATE/ACRE	NUMBER & TIMING OF APPLICATIONS
Alfalfa	2-6 pts/acre	Apply as soon as soil temperature reaches 55° at 6" and repeat after cutting as required.
Citrus	2-6 pts/acre	Multiple applications as required.
Cotton	3-6 pts/acre	2-3 applications from squaring to full boll development.
Cucurbit Vegetables such as: Cucumber, Cantaloupe, Watermelon, Summer and Winter Squash and Pumpkin.	2-6 pts/acre	2-3 applications from bloom through fruit sizing.
Leafy Vegetables and Brassica (Cole) such as: Broccoli, Cabbage, Cauliflower, Celery, Lettuce and Spinach.	2-4 pts/acre	2-4 applications from thinning to pre-harvest.
Root and Tuber Vegetables such as: Carrot, Potato, Onion, Garlic, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip and Yams.	2-5 pts/acre	2-3 applications during the season as required.
For aerial application use a maximum of 4 pints of FIRST CHOICE FOLIAR GOLD per 10 gallons of water		

NOTE: Rates suggested are for standard spray volumes of 20-150 gal/acre. For higher volume sprays, rate should be adjusted.

STORAGE: Store in cool dry area. Avoid extreme variations in temperature. A reversible separation of ingredients may occur after prolonged storage. This separation has no effect on quality or effectiveness of product and agitation will resuspend mixture.