



GUARANTEED ANALYSIS

Total Nitrogen (N).....6.0%
 2.4% Ammoniacal Nitrogen
 1.8% Nitrate Nitrogen
 1.8% Urea Nitrogen
 Available Phosphoric Acid (P₂O₅).....3.0%
 Soluble Potash (K₂O).....9.0%
 Sulfur (S).....4.0%
 Iron (Fe).....0.25%
 0.25% Chelated Iron
 Manganese (Mn).....0.10%
 0.10% Chelated Manganese
 Zinc (Zn).....0.10%
 0.10% Chelated Zinc

Nutrients derived from ammonium polyphosphate, ammonium thiosulfate, aqueous ammonia, citric acid, EDTA acid (Ethylenediaminetetraacetate), ferrous sulfate, iron EDTA, lo-biuret urea, manganese sulfate, potassium hydroxide, potassium nitrate, potassium phosphate, potassium thiosulfate, and zinc oxide.

Net weight: 10.38 pounds per gallon (1.25 kg/liter)
 NET CONTENTS _____ GALLONS(_____) LITER

DISTRIBUTED BY

Western Farm Service, Inc.
 P.O. Box 1168
 Fresno, CA 93715-1168

*GOLD'n GRO is a registered trademark of Itronics Metallurgical, Inc.
 FIRST CHOICE is a registered trademark of Western Farm Service, Inc.*

L03.10.05.32. OR

GOLD'n GRO 6-3-9 + 4%S

6-3-9 Liquid with 4% Sulfur and Chelated Iron, Chelated Manganese, Chelated Zinc

Directions for Use

This product is a chloride-free, lo-biuret fertilizer. GOLD'n GRO 6-3-9 + 4%S can be used on turf, ornamentals, wine, table, and raisin grapes, field crops including cereal grains, fruit and nut trees, citrus, and avocados. The total amount required per acre per year per crop may be determined by standard grower practice. Maximum quantity applied per acre per year should not exceed 60 gallons (623 pounds) in the State of Washington.

Spray applications for turf, ornamentals, all varieties of grapes, vegetables, other field crops including cereal grains, fruit and nut trees, citrus and avocados should be done using a mix no more concentrated than 2 quarts in 10 gallons of water. Do not foliar apply when temperature exceeds 90 degrees Fahrenheit.

Add the GOLD'n GRO products to water. Use a bottle test for mixtures with other products before proceeding with a full scale mix.

GOLD'n GRO 6-3-9 + 4%S may be applied through drip, sprinkler, and micro sprinkler irrigation or injected into the soil as in sidedress or listing operations. When GOLD'n GRO 6-3-9 + 4%S is injected in drip, drip tape or micro sprinklers, a jar test for compatibility with the irrigation water is recommended.

GOLD'n GRO 6-3-9 + 4%S is compatible with most fungicides, herbicides, and insecticides. When in doubt, do a jar test or contact your supplier. When combining with fungicides, herbicides, or insecticides, follow the manufacturer's instructions as to placement, dilution rates, and all other precautions.

Suggested Use Rates

Crop	Application Rate	Number and Timing of Applications
Alfalfa (Foliar)	1-8 qt./acre	Apply after each hay cutting as regrowth starts. On seed applications 4-6 times prior to harvest.
Alfalfa (Drench/Soil application)	1-12 gal./acre	Apply after each cutting.
Artichokes (Foliar)	1-2 qt./acre	Apply at bud stage, repeat beyond bud stage as needed.
Blackberries & Boysenberries, Raspberries, Strawberries (Foliar)	1-2 qt./acre	Apply pre-bloom, repeat as needed until berries color or soften.
Carrots (Foliar)	1-3 qt./acre	Begin applications when carrots are pencil size, repeat as needed.
Carrots (Drench/Soil application)	1-12 gal./acre	Begin applications at first irrigation (fertigation) or sidedress. Continue as needed.
Cotton (Foliar)	1-8 qt./acre	Begin applications at pin square. Repeat through the season as needed.
Citrus, Avocados (Foliar)	1-12 qt./acre	Application schedule will be dependent on crop, variety and cultural practices. Use higher rates on larger trees.
Citrus, Avocados (Drench/Soil application)	1-12 gal./acre	Begin applications at first flush of growth in spring. Repeat as necessary.
Deciduous fruit & nut trees (Foliar)	1-12 qt./acre	Begin applications after leaves appear. Application schedule will be dependant on crop, variety and cultural practices of the region.
Deciduous fruit and nut trees (Drench/Soil application)	1-12 gal./acre	Apply at or just before bud break. Repeat as necessary. Application schedule will be dependent on crop, variety and cultural practices of the region.
Grapes (Foliar)	1-8 qt./acre	Apply after leaves emerge. Repeat as necessary to veraison. Use lower rates early. Increase rates as foliage increases.
Grapes (Drench, Soil application)	1-12 gal./acre	Begin at or just before bud break. Repeat as necessary to veraison. A post-harvest application is recommended.
Potatoes (Foliar)	1-8 qt./acre	Apply at bud formation; repeat as necessary.
Potatoes (Drench/Soil application)	1-12 gal./acre	Begin applications in the first irrigation (fertigation) repeat as necessary.
Corn (Foliar)	1-8 qt./acre	Apply at 6" height and repeat as needed.
Nursery, Greenhouse, Ornamentals (Drench)	16-48 oz./100 gal.	Use in every irrigation. Drench to soak planting medium.
Field and vegetable crops, Not otherwise listed (Foliar)	1-3 qt./acre	Begin applications when plants are 6 to 8 inches tall (6 to 8 inches across for low-growing plants). Continue as necessary. Application schedule will vary depending on crop, variety and region.
Field and vegetable crops, Not otherwise listed (Drench/Soil application)	1-12 gal./acre	Begin applications at first sprinkler (fertigation) or sidedress application after emergence. Application schedule will be dependent on crop, variety and cultural practices of the region.
General use (Foliar application)	1-12 qt./acre	Application schedule will be dependent on crop, variety and cultural practices of the region.
General use (Drench/Soil application)	1-12 gal./acre	See directions for use concerning application methods.

Material of Construction

The concentrated product is corrosive for iron, copper, zinc, brass, bronze and aluminum. Tanks for storage of concentrated product can be polypropylene, 304 or 316 stainless steel, or cross-linked polyethylene. Fiberglass, iron, and aluminum tanks should be avoided. It is okay to use fiberglass and 304 stainless spray tanks for ground spray rigs and aluminum for aerial applications. Do not store fertilizer mix in spray tanks. Flush lines after using.

CONDITIONS OF SALE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, damage caused by drift to other plants or crops, and personal injury.

CAUTION: KEEP OUT REACH OF CHILDREN. Not for internal consumption. Avoid contact. A slight sulfur odor may occur for a few hours after spray application.

WARNING: This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

NOTE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Information regarding the contents and levels of metals in this product is available on the internet at www.aapfco.org/metals.htm