



FIRST CHOICE[®]

Bean Chaser™

10-0-0

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

GUARANTEED ANALYSIS:

Total Nitrogen (N)	10.0%
0.55% Ammoniacal Nitrogen	
9.05% Nitrate Nitrogen	
0.40% Urea Nitrogen	
Calcium (Ca)	10.60%
Magnesium (Mg)	1.20%
Boron (B)	0.05%
Copper (Cu)	0.05%
0.05% Chelated Copper	
Iron (Fe)	0.10%
0.10% Chelated Iron	
Manganese (Mn)	0.10%
0.10% Chelated Manganese	
Molybdenum (Mo)	0.001%
Zinc (Zn)	0.05%
0.05% Chelated Zinc	

Derived from Calcium Nitrate, Calcium Ammonium Nitrate, Urea, Magnesium Nitrate, Boric Acid, Copper Ethylenediaminetetraacetate (EDTA), Iron EDTA, Manganese EDTA, Sodium Molybdate and Zinc EDTA.

GENERAL INFORMATION

FIRST CHOICE BEAN CHASER is a water-soluble foliar nutritional fertilizer of macro- and micro nutrients. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. Foliar fertilization is a supplement to a regular fertilizer program and will not supply the total nutrients required by a crop. FIRST CHOICE BEAN CHASER is recommended as a foliar nutrition to be applied to crops where disorders due to insufficient calcium are anticipated.

FOR COMMERCIAL USE ONLY

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use, storage or handling of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. These risks include, but are not limited to, damage to plants, crops and animals to which the material is applied, damage caused by drift to other plants or crops, and personal injury.

CAUTION
Keep Out of Reach of Children

The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

WARNING: This fertilizing material contains Boron (B). Use on any crops other than those recommended may result in serious injury to the crop(s).

NET CONTENTS: 2.5 GALLONS (9.463 LITERS)
13.0 lbs./gallon (1.56 kgs./liter)

Manufactured For:
Western Farm Service, Inc.
P. O. Box 1168
Fresno California 93715

FIRST CHOICE is a registered trademark of Western Farm Service, Inc.
BEAN CHASER is a trademark of Western Farm Service, Inc.
BREAKTHRU is a registered trademark of Goldschmidt Chemical Corporation

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

DIRECTIONS FOR USE

Dissolve the desired amount of FIRST CHOICE BEAN CHASER in water before adding to the spray tank. With proper agitation, FIRST CHOICE BEAN CHASER can be added into the spray tank with agitation running. Do not store solution mixed with pesticide! Spray immediately after mixing. Add FIRST CHOICE BEAN CHASER to the spray tank before adding pesticides.

RECOMMENDATIONS FOR USE

CROP	RATE/ACRE	NUMBER & TIMING OF APPLICATIONS
Beans	2 pints/acre 1 st application	1 st application between 2 nd and 3 rd trifoliolate leaf stage.
	2 pints/acre 2 nd application	2 nd application at 1 st bloom stage (less than 10% bloom).

Rates suggested are for standard spray volumes of 40 – 150 gallons per acre. For higher volume sprays, rate should be adjusted. Caution: Store FIRST CHOICE BEAN CHASER in temperatures between 41°F to 104°F and avoid extreme variations in temperature. A reversible separation of ingredients may occur after long storage, this however does not affect quality or effectiveness of the product.

When applying to beans, use a high analysis organo silicone such as First Choice BreakThru® or a high analysis non-ionic surfactant such as First Choice A Plus Spreader during the recommended timings listed above. Avoid applications with the organo silicone when temperatures are elevated >90°F or irrigations are to be applied within 4 hours of application. Maximum effectiveness will be obtained when applied early in the morning or after dusk.